

Contribution to the 18th International IUPAP Conference on Few-Body Problems in Physics (FB18), Santos, Brasil, August 21-26, 2006

LPNHE 2006-13

Strong and Weak Interactions in $B \rightarrow \pi^+\pi^-K$ Decays

B. Loiseau^{a *}, B. El-Bennich^a, A. Furman^b, R. Kamiński^c, L. Leśniak^c

^aLPNHE (IN2P3–CNRS–Universités Paris 6 et 7), Groupe Théorie,
Université Pierre et Marie Curie, 4 place Jussieu, 75252 Paris, France

^bul. Bronowicka 85/26, 30-091 Kraków, Poland

^cDivision of Theoretical Physics, The Henryk Niewodniczański Institute of Nuclear
Physics, Polish Academy of Sciences, 31-342 Kraków, Poland

To describe the weak three-body decays $B \rightarrow \pi^+\pi^-K$, we recently derived amplitudes based on two-body QCD factorization followed by $\pi^+\pi^-$ final state interactions in isoscalar S - and isovector P -waves. We study here the sensitivity of the results to the values of the B to $f_0(980)$ transition form factor and to the effective decay constant of the $f_0(980)$.

1. INTRODUCTION

It is important to understand charmless three-body B decays to probe the standard model. These decays are sensitive to CP violation and supply information on strong interactions. To interpret in a reliable way weak decay observables it is important to take into account final state interactions between produced meson pairs. In the weak decays $B \rightarrow \pi^+\pi^-K$ [1, 2] one sees maxima around the $\pi\pi$ effective mass distributions in the $\rho(770)^0$ and $f_0(980)$ resonance regions.

For $\pi\pi$ effective mass $m_{\pi\pi}$ up to 1.2 GeV, the contribution of the isospin-zero S -wave $(\pi^+\pi^-)_S$ final state interactions was described in Ref. [3] and that of the isospin-one P wave $(\pi^+\pi^-)_P$ was included in Ref. [4]. The amplitudes, based on the QCD factorization approach without the inclusion of hard-spectator and annihilation terms, underestimate the B to $\rho(770)^0K$ and $f_0(980)K$ branching fractions. Therefore, phenomenological amplitudes arising from enhanced $c\bar{c}$ loop diagrams (charming penguin terms [5]) were added. Our presentation at the conference was based on the work described in Ref. [4]. Here we show the sensitivity of the model [4] to two inputs of the S -wave amplitude: the B to $f_0(980)$ transition form factor and the effective decay constant of the $f_0(980)$.

2. WEAK DECAY AMPLITUDES FOR $B \rightarrow \pi^+\pi^-K$

The amplitudes for the weak decays $B \rightarrow (\pi^+\pi^-)_{S(P)}K$ are derived [3, 4] in the QCD factorization framework [6, 7]. As a first approximation, corrections arising from annihilation topologies and hard gluon scattering with the spectator quark are not included. These also contain several phenomenological parameters (see for instance [7]).

*B. E. and B.L. acknowledge useful discussions with J. P. Dedonder, O. Leitner and Craig Roberts.




Figure 1. Quark-line diagrams for two-body B^- decay into K^- and a $\pi\pi$ state $R_S(m_{\pi\pi})$ or $R_P(m_{\pi\pi})$ in an isoscalar S - or isovector P -wave, respectively. The filled circle represents the weak and electroweak decays via tree or penguin type diagrams.

For the $B \rightarrow (\pi^+\pi^-)_S K$ decay amplitudes, we consider the three-body $\pi^+\pi^-K$ final state as arising from a quasi two-body one with the produced $\pi^+\pi^-$ pair being in an isospin-0 S -wave state R_S of mass $m_{\pi\pi}$. For $m_{\pi\pi} = m_{f_0}$ (mass of the $f_0(980)$) this R_S state is the $f_0(980)$ resonance. Our amplitudes have a weak two-body decay part based on operator product expansion, heavy quark limit and QCD factorization [6] followed by the strong decay of R_S into $(\pi^+\pi^-)_S$ with inclusion of rescattering. One has [3, 4],

$$\langle (\pi^+\pi^-)_S K^- | H | B^- \rangle = M_K^S(m_{\pi\pi}) + M_{R_S}^S(m_{\pi\pi}) , \quad (1)$$

$$M_K^S(m_{\pi\pi}) = \frac{G_F}{\sqrt{2}} f_K (M_B^2 - m_{\pi\pi}^2) F_0^{B \rightarrow R_S}(M_K^2) \left\{ \lambda_u [a_1 + a_4^u - a_4^c + (a_6^c - a_6^u) r - S_u] \right. \\ \left. + \lambda_t (a_6^c r - a_4^c - S_t) \right\} \sqrt{\frac{2}{3}} \Gamma_{R_S \pi\pi}^n(m_{\pi\pi}) , \quad (2)$$

$$M_{R_S}^S(m_{\pi\pi}) = \frac{G_F}{\sqrt{2}} (M_B^2 - M_K^2) \left\{ \langle R_S | \bar{s}s | 0 \rangle \frac{2F_0^{B \rightarrow K}(m_{\pi\pi}^2)}{m_b - m_s} [\lambda_u (a_6^c - a_6^u) + \lambda_t a_6^c] \right. \\ \left. - N_K (\lambda_u S_u + \lambda_t S_t) \right\} \sqrt{\frac{2}{3}} \Gamma_{R_S \pi\pi}^s(m_{\pi\pi}) . \quad (3)$$

In Eqs. (2) and (3), G_F is the Fermi constant, f_K the kaon decay constant and M_B , M_K , m_b and m_s the B -meson, kaon, b - and s -quark masses, respectively. The functions $F_0^{B \rightarrow R_S}(M_K^2)$ and $F_0^{B \rightarrow K}(m_{\pi\pi}^2)$ are the B^- to R_S or K^- transition form factors. The $\lambda_{u,t}$ are products of the CKM matrix elements, $\lambda_u = V_{ub}V_{us}^*$ and $\lambda_t = V_{tb}V_{ts}^*$. The a_i are the scale dependent effective coefficients built from the Wilson coefficients and including next-to-leading-order QCD corrections [6, 7]. The chiral factor $r = 2M_K^2/[(m_b+m_u)(m_s+m_u)]$, m_u being the u -quark mass. The phenomenological charming penguin parameters S_u and S_t are added to the QCD factorization terms. In Eq. (3) the weight factor N_K is chosen to be $N_K = f_K F_0^{B \rightarrow R_S}(M_K^2)$ as in Eq. (2).

The amplitude $M_K^S(m_{\pi\pi})$ matches the topology of Figure 1 (a) with the production of a K^- meson from the vacuum plus a B^- transition into an $R_S(m_{\pi\pi})$ state. The amplitude $M_{R_S}^S(m_{\pi\pi})$ corresponds to the topology of Figure 1(b) with the emission of an $R_S(m_{\pi\pi})$ state from the vacuum plus a B^- transition into a K^- meson.

In Eqs. (2) and (3), the non-strange $\Gamma_{R_S\pi\pi}^n(m_{\pi\pi})$ and strange $\Gamma_{R_S\pi\pi}^s(m_{\pi\pi})$ vertex functions describe the strong decay of the state $R_S(m_{\pi\pi})$ into two pions and include $\pi^+\pi^-$ rescattering [3]. One can write

$$\langle(\pi\pi)_S|\bar{s}s|0\rangle = \Gamma_{R_S\pi\pi}^s(m_{\pi\pi})\langle R_S|\bar{s}s|0\rangle = \sqrt{2}B_0\Gamma_1^{s*}(m_{\pi\pi}) \quad (4)$$

where $\Gamma_1^s(m_{\pi\pi}) = \langle 0|\bar{s}s|(\pi\pi)_S\rangle/(\sqrt{2}B_0)$ is the strange scalar form factor and the normalization constant $B_0 = -\langle 0|\bar{q}q|0\rangle/f_\pi^2$, f_π being the pion decay constant. Replacing $\bar{s}s$ by $\bar{n}n = (\bar{u}u + \bar{d}d)/\sqrt{2}$ in Eq. (4) gives the non-strange vertex function $\Gamma_{R_S\pi\pi}^n(m_{\pi\pi})$ in terms of the non-strange scalar form factor $\Gamma_1^n(m_{\pi\pi})$. Defining a scalar decay constant $f_{R_S}^s$ by

$$\langle R_S|\bar{s}s|0\rangle = m_{R_S}f_{R_S}^s, \quad (5)$$

one then obtains from Eq. (4),

$$\Gamma_{R_S\pi\pi}^s(m_{\pi\pi}) = \chi\Gamma_1^{s*}(m_{\pi\pi}) \text{ with } \chi = \sqrt{2}B_0/(m_{R_S}f_{R_S}^s). \quad (6)$$

If we assume $\Gamma_{R_S\pi\pi}^n(m_{\pi\pi}) = \chi\Gamma_1^{n*}(m_{\pi\pi})$ and identify $R_S(m_{f_0})$ with $f_0(980)$, we normalize χ by [3]

$$\chi = g_{f_0\pi\pi}/[m_{f_0}\Gamma_{\text{tot}}(f_0)|\Gamma_1^n(m_{f_0})|], \quad (7)$$

where $\Gamma_{\text{tot}}(f_0)$ is the total $f_0(980)$ width.

Note that replacing $\sqrt{2/3}\Gamma_{R_S\pi\pi}^{n,s}(m_{\pi\pi})$ by 1 in Eqs. (2) and (3) leads to a two-body $B^- \rightarrow R_S K^-$ decay amplitude.

In the $B \rightarrow (\pi^+\pi^-)_P K$ decay amplitudes, the produced $\pi^+\pi^-$ pair is in an isovector P -wave $(\pi^+\pi^-)_P$ state $R_P(m_{\pi\pi})$ identified as the $\rho(770)^0$ resonance. The explicit expression of the $B^- \rightarrow (\pi^+\pi^-)_P K^-$ amplitude is given in Ref. [4]. As the amplitudes underestimate the B to $\rho(770)^0 K$ branching fraction, we also introduce a phenomenological charming penguin term depending on two complex parameters proportional to λ_u and λ_t .

The complete $B^- \rightarrow (\pi^+\pi^-)_{S+P} K^-$ amplitude is obtained by adding the S -wave amplitude of Eq. (1) to that of the P -wave [4]. Replacing λ_u, λ_t by λ_u^*, λ_t^* gives the $B^+ \rightarrow (\pi^+\pi^-)_S K^+$ amplitude. The expressions for the neutral B -decay and $B^+ \rightarrow (\pi^+\pi^-)_P K^+$ amplitudes can be found in [3] and [4]. In the results shown below we use the same input parameters as in Ref. [4] unless otherwise stated.

3. FIT, RESULTS, DISCUSSIONS AND CONCLUSIONS

In Ref. [4], we use $F_0^{B \rightarrow (\pi\pi)S}(M_K^2) = 0.46$ although recent calculations [8, 9, 10] give a value close to 0.25. Furthermore, the $\Gamma_1^{n,s}(m_{\pi\pi})$ depend on some poorly determined low energy constants of chiral perturbation theory. Using their latest determinations [11], the moduli of the $\Gamma_1^{n,s}(m_{\pi\pi})$ are larger by a factor of 1.25 than those of [4] in the $f_0(980)$ range. However, the constant value of $\chi|\Gamma_1^n(m_{f_0})|$ (see Eq. (7)) limits the

sensitivity to this variation. With these new inputs for the S -wave, the fit to the Belle and BaBar collaboration data (see references in [4]) is similar to that of Ref. [4]. The $\chi^2/d.o.f. = 346/(222 - 8) = 1.62$ in model [4] and 1.65 here. In this new fit, the resulting charming penguin parameters are modified and in particular, to compensate the decrease of $F_0^{B \rightarrow (\pi\pi)S}(M_K^2)$ from 0.46 to 0.25, the modulus of S_u increases.




Figure 2. The $m_{\pi\pi}$ distributions in $B^\pm \rightarrow \pi^+\pi^-K^\pm$ decays (data from Ref. [1]). The solid line results from the fit with $F_0^{B \rightarrow (\pi\pi)S}(M_K^2) = 0.25$ and with the $\Gamma_1^{n,s}(m_{\pi\pi})$ of larger moduli in the $f_0(980)$ range (see text). The vertical lines delimit the region of the fit.

In Fig. 2 the $m_{\pi\pi}$ branching fraction distributions are compared to the Belle data [1]. One sees an asymmetry in the number of events between the $B^- \rightarrow \pi^+\pi^-K^-$ and $B^+ \rightarrow \pi^+\pi^-K^+$ decays for the $\rho(770)^0$ and $f_0(980)$ regions. This results in a large direct CP asymmetry for $B^\pm \rightarrow \rho(770)^0K^\pm$ decays of 0.32 ± 0.03 to be compared with 0.30 ± 0.14 and 0.32 ± 0.16 from Belle and BaBar collaborations, Refs. [1] and [4] in [4], respectively.

It was found in Ref. [8] that the experimental average branching fractions of two-body $B^\pm \rightarrow f_0(980)K^\pm$ decays, $\mathcal{B}_{f_0}$, could be reproduced without charming penguin terms. However, we have important differences with Ref. [8] in the S -wave inputs.

First, concerning the $M_{R_S}^S(m_{\pi\pi})$ amplitude, we have shown in section 2 that, with $R_S \equiv f_0$, we use an effective scalar decay constant $f_{f_0}^s = \sqrt{2}B_0/(m_{f_0}\chi) = 94$ MeV with the input parameters of Ref. [4] or 117.5 MeV with the new $\Gamma_1^{n,s}$ just considered above. These values are to be compared with those of 370 MeV determined from QCD sum rules in Ref. [8] or of 245 MeV obtained in Ref. [12] applying the Dyson-Schwinger equations, which respect dynamical chiral symmetry breaking in modeling scalar mesons.

With $F_0^{B \rightarrow (\pi\pi)S}(M_K^2) = 0.46$, if we set $S_u = S_t = 0$ and $\Gamma_1^{s*}(m_{\pi\pi}) = 0$ (equivalent to an effective $f_{f_0}^s = 0$), $\mathcal{B}_{f_0} = 2.19 (0.67) \times 10^{-6}$ when integrating the $m_{\pi\pi}$ average distribution from 0.900 to 1.060 GeV. Here and below, the cited number in parenthesis corresponds to

the fit with $F_0^{B \rightarrow (\pi\pi)S}(M_K^2) = 0.25$ and with the $\Gamma_1^{n,s}(m_{\pi\pi})$ of larger moduli in the $f_0(980)$ range. With the addition of $M_{R_S}^S(m_{\pi\pi})$ of [4] (effective $f_{f_0}^s = 94$ (117.5) MeV), $\mathcal{B}_{f_0} = 2.66$ (1.09) $\times 10^{-6}$. If we multiply $\Gamma_1^{s*}(m_{\pi\pi})$ by 4 (3.15) (effective $f_{f_0}^s = 376$ (370) MeV), $\mathcal{B}_{f_0} = 4.50$ (2.49) $\times 10^{-6}$. Remind that the corresponding Belle value is $6.06 \pm 1.08 \times 10^{-6}$. In our case, if we add the contributions of S_u and S_t of Ref. [4], then $\mathcal{B}_{f_0}$ increases from 2.66 (1.09) $\times 10^{-6}$ to 6.93 (6.59) $\times 10^{-6}$.

Secondly, the fit to experimental data of Ref. [8] includes hard spectator scattering terms with the parameters X_A (plus annihilation terms with the parameters X_H). As stated in Ref. [13], the $a_1(f_0 K)$ receives a large contribution from hard spectator interaction which will enhance the $M_K^S(m_{f_0})$ of Ref. [8].

In summary, uncertainties in $F_0^{B \rightarrow (\pi\pi)S}(M_K^2)$ and in the S -wave scalar form factors lead to variations of charming penguin parameters, in particular of S_u . The scalar form factors that we use give low values for the effective $f_{f_0}^s$ decay constant equal to 94 or 117.5 MeV, to be compared to 370 MeV [8] or 245 MeV [12]. Despite these uncertainties, our conclusions [4] are unchanged. Our theoretical model is based on quasi two-body QCD factorization followed by S - and P -wave final state interactions between the produced $\pi\pi$ pairs. These interactions are constrained by other experiments, unitarity and chiral symmetry. Our model gives a good fit of the three-body $B \rightarrow \pi^+\pi^-K$ decay data. In particular it describes well the interference between the $f_0(980)$ and $\rho(770)^0$ resonances.

REFERENCES

1. K. Abe *et al.*, Belle Collaboration, hep-ex/0509001, *Search for direct CP violation in three body charmless $B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$ decays*.
2. B. Aubert, *et al.*, BaBar Collaboration, Phys. Rev. D **73** (2006) 031101(R), *Measurements of neutral B decay branching fractions to $K_S^0 \pi^+ \pi^-$ final states and the charge asymmetry of $B^0 \rightarrow K^{*+} \pi^-$* .
3. A. Furman, R. Kamiński, L. Leśniak and B. Loiseau, Phys. Lett. B **622** (2005) 207, *Long-distance effects and final state interactions in $B \rightarrow \pi\pi K$ and $B \rightarrow K\bar{K}K$ decays*.
4. B. El-Bennich, A. Furman, R. Kamiński, L. Leśniak and B. Loiseau, hep-ph/0608205, *Interference between $f_0(980)$ and $\rho(770)^0$ resonances in $B \rightarrow \pi^+\pi^-K$ decays*.
5. M. Ciuchini, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. B **501** (1997) 271, *Charming penguins in B decays*.
6. G. Buchalla, A. J. Buras, M. E. Lautenbacher, Review of Modern Physics **68** (1996) 1125, *Weak decays beyond leading logarithms*.
7. M. Beneke and M. Neubert, Nucl. Phys. B **675** (2003) 333, *QCD factorization for $B \rightarrow PP$ and $B \rightarrow PV$ decays*.
8. H. Y. Cheng, C. K. Chua and K. C. Yang, Phys. Rev. D **73** (2006) 014017, *Charmless hadronic B decays involving scalar mesons: Implications on the nature of light scalar mesons*.
9. O. M. A. Leitner, B. El-Bennich, B. Loiseau and J.-P. Dedonder, hep-ph/0609062, *Pseudoscalar - scalar transition form factors in covariant light front dynamics*.
10. B. El-Bennich, O. M. A. Leitner, B. Loiseau and J.-P. Dedonder, hep-ph/0609154, *Form factors in $B \rightarrow f_0(980)$ and $D \rightarrow f_0(980)$ transitions from dispersion relations*; Idem, in these proceedings, B. El-Bennich's talk, hep-ph/0610229, *Scalar meson prop-*

erties from D-meson decays.

11. T. A. Lähde and U. G. Meißner, Phys. Rev. D **74** (2006) 034021, *Improved Analysis of J/ψ Decays into a Vector Meson and Two Pseudoscalars.*
12. M. S. Bhagwat, A. Höll, A. Krassnigg, C. D. Roberts, in these proceedings, *Theory and Phenomenology of Hadrons.*
13. H. Y. Cheng and K. C. Yang, Phys. Rev. D **71** (2005) 054020, *$B \rightarrow f_0(980)K$ decays and subleading corrections.*